

# Sample Spaces, Events, Probability

CS 3130/ECE 3530:  
Probability and Statistics for Engineers

Jan 12, 2023

# Sets

## Definition

A **set** is a collection of unique objects.

# Sets

## Definition

A **set** is a collection of unique objects.

Here “objects” can be concrete things (people in class, schools in PAC-12), or abstract things (numbers, colors).

# Sets

## Definition

A **set** is a collection of unique objects.

Here “objects” can be concrete things (people in class, schools in PAC-12), or abstract things (numbers, colors).

Examples:

$$A = \{3, 8, 31\}$$

# Sets

## Definition

A **set** is a collection of unique objects.

Here “objects” can be concrete things (people in class, schools in PAC-12), or abstract things (numbers, colors).

Examples:

$$A = \{3, 8, 31\}$$

$$B = \{\text{apple, pear, orange, grape}\}$$

# Sets

## Definition

A **set** is a collection of unique objects.

Here “objects” can be concrete things (people in class, schools in PAC-12), or abstract things (numbers, colors).

Examples:

$$A = \{3, 8, 31\}$$

$$B = \{\text{apple, pear, orange, grape}\}$$

**Not** a valid set definition:  $C = \{1, 2, 3, 4, 2\}$

# Sets

- ▶ Order in a set does not matter!

$$\{1, 2, 3\} = \{3, 1, 2\} = \{1, 3, 2\}$$

# Sets

- ▶ Order in a set does not matter!

$$\{1, 2, 3\} = \{3, 1, 2\} = \{1, 3, 2\}$$

- ▶ When  $x$  is an element of  $A$ , we denote this by:

$$x \in A.$$



# Sets

- ▶ Order in a set does not matter!

$$\{1, 2, 3\} = \{3, 1, 2\} = \{1, 3, 2\}$$

- ▶ When  $x$  is an element of  $A$ , we denote this by:

$$x \in A.$$

- ▶ If  $x$  is not in a set  $A$ , we denote this as:

$$x \notin A.$$

# Sets

- ▶ Order in a set does not matter!

$$\{1, 2, 3\} = \{3, 1, 2\} = \{1, 3, 2\}$$

- ▶ When  $x$  is an element of  $A$ , we denote this by:

$$x \in A.$$

- ▶ If  $x$  is not in a set  $A$ , we denote this as:

$$x \notin A.$$

- ▶ The “empty” or “null” set has no elements:

$$\emptyset = \{ \}$$

# Some Important Sets

- ▶ Integers:

$$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$$

# Some Important Sets

- ▶ Integers:

$$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$$

- ▶ Natural Numbers:

$$\mathbb{N} = \{0, 1, 2, 3, \dots\}$$

# Some Important Sets

- ▶ Integers:

$$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$$

- ▶ Natural Numbers:

$$\mathbb{N} = \{0, 1, 2, 3, \dots\}$$

- ▶ Real Numbers:

$\mathbb{R}$  = “any number that can be written in decimal form”

# Some Important Sets

- ▶ Integers:

$$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$$

- ▶ Natural Numbers:

$$\mathbb{N} = \{0, 1, 2, 3, \dots\}$$

- ▶ Real Numbers:

$\mathbb{R}$  = “any number that can be written in decimal form”

$$5 \in \mathbb{R}, \quad 17.42 \in \mathbb{R}, \quad \pi = 3.14159\dots \in \mathbb{R}$$

# Building Sets Using Conditionals

# Building Sets Using Conditionals

- ▶ Alternate way to define natural numbers:

$$\mathbb{N} = \{x \in \mathbb{Z} : x \geq 0\}$$



# Building Sets Using Conditionals

- ▶ Alternate way to define natural numbers:

$$\mathbb{N} = \{x \in \mathbb{Z} : x \geq 0\}$$

- ▶ Set of even integers:

$$\{x \in \mathbb{Z} : x \text{ is divisible by } 2\}$$

# Building Sets Using Conditionals

- ▶ Alternate way to define natural numbers:

$$\mathbb{N} = \{x \in \mathbb{Z} : x \geq 0\}$$

- ▶ Set of even integers:

$$\{x \in \mathbb{Z} : x \text{ is divisible by } 2\}$$

- ▶ Rationals:

$$\mathbb{Q} = \{p/q : p, q \in \mathbb{Z}, q \neq 0\}$$

# Subsets

## Definition

A set  $A$  is a **subset** of another set  $B$  if every element of  $A$  is also an element of  $B$ , and we denote this as  $A \subseteq B$ .

# Subsets

## Definition

A set  $A$  is a **subset** of another set  $B$  if every element of  $A$  is also an element of  $B$ , and we denote this as  $A \subseteq B$ .

Examples:

# Subsets

## Definition

A set  $A$  is a **subset** of another set  $B$  if every element of  $A$  is also an element of  $B$ , and we denote this as  $A \subseteq B$ .

Examples:

- ▶  $\{1, 9\} \subseteq \{1, 3, 9, 11\}$

# Subsets

## Definition

A set  $A$  is a **subset** of another set  $B$  if every element of  $A$  is also an element of  $B$ , and we denote this as  $A \subseteq B$ .

Examples:

- ▶  $\{1, 9\} \subseteq \{1, 3, 9, 11\}$
- ▶  $\mathbb{Q} \subseteq \mathbb{R}$

# Subsets

## Definition

A set  $A$  is a **subset** of another set  $B$  if every element of  $A$  is also an element of  $B$ , and we denote this as  $A \subseteq B$ .

Examples:

- ▶  $\{1, 9\} \subseteq \{1, 3, 9, 11\}$
- ▶  $\mathbb{Q} \subseteq \mathbb{R}$
- ▶  $\{\text{apple, pear}\} \not\subseteq \{\text{apple, orange, banana}\}$

# Subsets

## Definition

A set  $A$  is a **subset** of another set  $B$  if every element of  $A$  is also an element of  $B$ , and we denote this as  $A \subseteq B$ .

Examples:

- ▶  $\{1, 9\} \subseteq \{1, 3, 9, 11\}$
- ▶  $\mathbb{Q} \subseteq \mathbb{R}$
- ▶  $\{\text{apple, pear}\} \not\subseteq \{\text{apple, orange, banana}\}$
- ▶  $\emptyset \subseteq A$  for any set  $A$



# Subsets

## Definition

A set  $A$  is a **subset** of another set  $B$  if every element of  $A$  is also an element of  $B$ , and we denote this as  $A \subseteq B$ .

Examples:

- ▶  $\{1, 9\} \subseteq \{1, 3, 9, 11\}$
- ▶  $\mathbb{Q} \subseteq \mathbb{R}$
- ▶  $\{\text{apple, pear}\} \not\subseteq \{\text{apple, orange, banana}\}$
- ▶  $\emptyset \subseteq A$  for any set  $A$
- ▶  $A \subseteq A$  for any set  $A$  (but  $A \not\subset A$ )

# Sample Spaces

## Definition

A **sample space** is the set of all possible outcomes of an experiment. We'll denote a sample space as  $\Omega$ .

# Sample Spaces

## Definition

A **sample space** is the set of all possible outcomes of an experiment. We'll denote a sample space as  $\Omega$ .

Examples:

- ▶ Coin flip:  $\Omega = \{H, T\}$

# Sample Spaces

## Definition

A **sample space** is the set of all possible outcomes of an experiment. We'll denote a sample space as  $\Omega$ .

Examples:

- ▶ Coin flip:  $\Omega = \{H, T\}$
- ▶ Roll a 6-sided die:  $\Omega = \{1, 2, 3, 4, 5, 6\}$

# Sample Spaces

## Definition

A **sample space** is the set of all possible outcomes of an experiment. We'll denote a sample space as  $\Omega$ .

Examples:

- ▶ Coin flip:  $\Omega = \{H, T\}$
- ▶ Roll a 6-sided die:  $\Omega = \{1, 2, 3, 4, 5, 6\}$
- ▶ Pick a ball from a bucket of red/black balls:  
 $\Omega = \{R, B\}$

# Sample Spaces

## Definition

A **sample space** is the set of all possible outcomes of an experiment. We'll denote a sample space as  $\Omega$ .

Examples:

- ▶ Coin flip:  $\Omega = \{H, T\}$
- ▶ Roll a 6-sided die:  $\Omega = \{1, 2, 3, 4, 5, 6\}$
- ▶ Pick a ball from a bucket of red/black balls:  
 $\Omega = \{R, B\}$
- ▶ Tossing 2 coins?

# Sample Spaces

## Definition

A **sample space** is the set of all possible outcomes of an experiment. We'll denote a sample space as  $\Omega$ .

Examples:

- ▶ Coin flip:  $\Omega = \{H, T\}$
- ▶ Roll a 6-sided die:  $\Omega = \{1, 2, 3, 4, 5, 6\}$
- ▶ Pick a ball from a bucket of red/black balls:  
 $\Omega = \{R, B\}$
- ▶ Tossing 2 coins?
- ▶ Shuffling deck of 52 cards?

# Events

## Definition

An **event** is a subset of a sample space.



# Events

## Definition

An **event** is a subset of a sample space.

Examples:

- ▶ You roll a die and get an even number:  
 $\{2, 4, 6\} \subseteq \{1, 2, 3, 4, 5, 6\}$

# Events

## Definition

An **event** is a subset of a sample space.

Examples:

- ▶ You roll a die and get an even number:  
 $\{2, 4, 6\} \subseteq \{1, 2, 3, 4, 5, 6\}$
- ▶ You flip a coin and it comes up “heads”:  
 $\{H\} \subseteq \{H, T\}$

# Events

## Definition

An **event** is a subset of a sample space.

Examples:

- ▶ You roll a die and get an even number:  
 $\{2, 4, 6\} \subseteq \{1, 2, 3, 4, 5, 6\}$
- ▶ You flip a coin and it comes up “heads”:  
 $\{H\} \subseteq \{H, T\}$
- ▶ Your code takes longer than 5 seconds to run:  
 $(5, \infty) \subseteq \mathbb{R}$

# Set Operations: Union

## Definition

The **union** of two sets  $A$  and  $B$ , denoted  $A \cup B$  is the set of all elements in either  $A$  or  $B$  (or both).

# Set Operations: Union

## Definition

The **union** of two sets  $A$  and  $B$ , denoted  $A \cup B$  is the set of all elements in either  $A$  or  $B$  (or both).

When  $A$  and  $B$  are events,  $A \cup B$  means that event  $A$  or event  $B$  happens (or both).

# Set Operations: Union

## Definition

The **union** of two sets  $A$  and  $B$ , denoted  $A \cup B$  is the set of all elements in either  $A$  or  $B$  (or both).

When  $A$  and  $B$  are events,  $A \cup B$  means that event  $A$  or event  $B$  happens (or both).

Example:

$A = \{1, 3, 5\}$     “an odd roll”

$B = \{1, 2, 3\}$     “a roll of 3 or less”

# Set Operations: Union

## Definition

The **union** of two sets  $A$  and  $B$ , denoted  $A \cup B$  is the set of all elements in either  $A$  or  $B$  (or both).

When  $A$  and  $B$  are events,  $A \cup B$  means that event  $A$  or event  $B$  happens (or both).

Example:

$$A = \{1, 3, 5\} \quad \text{"an odd roll"}$$

$$B = \{1, 2, 3\} \quad \text{"a roll of 3 or less"}$$

$$A \cup B = \{1, 2, 3, 5\}$$

# Set Operations: Intersection

## Definition

The **intersection** of two sets  $A$  and  $B$ , denoted  $A \cap B$  is the set of all elements in both  $A$  and  $B$ .



# Set Operations: Intersection

## Definition

The **intersection** of two sets  $A$  and  $B$ , denoted  $A \cap B$  is the set of all elements in both  $A$  and  $B$ .

When  $A$  and  $B$  are events,  $A \cap B$  means that both event  $A$  and event  $B$  happen.

# Set Operations: Intersection

## Definition

The **intersection** of two sets  $A$  and  $B$ , denoted  $A \cap B$  is the set of all elements in both  $A$  and  $B$ .

When  $A$  and  $B$  are events,  $A \cap B$  means that both event  $A$  and event  $B$  happen.

Example:

$A = \{1, 3, 5\}$     “an odd roll”

$B = \{1, 2, 3\}$     “a roll of 3 or less”

# Set Operations: Intersection

## Definition

The **intersection** of two sets  $A$  and  $B$ , denoted  $A \cap B$  is the set of all elements in both  $A$  and  $B$ .

When  $A$  and  $B$  are events,  $A \cap B$  means that both event  $A$  and event  $B$  happen.

Example:

$$A = \{1, 3, 5\} \quad \text{"an odd roll"}$$

$$B = \{1, 2, 3\} \quad \text{"a roll of 3 or less"}$$

$$A \cap B = \{1, 3\}$$

# Set Operations: Intersection

## Definition

The **intersection** of two sets  $A$  and  $B$ , denoted  $A \cap B$  is the set of all elements in both  $A$  and  $B$ .

When  $A$  and  $B$  are events,  $A \cap B$  means that both event  $A$  and event  $B$  happen.

Example:

$$A = \{1, 3, 5\} \quad \text{"an odd roll"}$$

$$B = \{1, 2, 3\} \quad \text{"a roll of 3 or less"}$$

$$A \cap B = \{1, 3\}$$

Note: If  $A \cap B = \emptyset$ , we say  $A$  and  $B$  are **disjoint**.

# Set Operations: Complement

## Definition

The **complement** of a set  $A \subseteq \Omega$ , denoted  $A^c$ , is the set of all elements in  $\Omega$  that are not in  $A$ .

# Set Operations: Complement

## Definition

The **complement** of a set  $A \subseteq \Omega$ , denoted  $A^c$ , is the set of all elements in  $\Omega$  that are not in  $A$ .

When  $A$  is an event,  $A^c$  means that the event  $A$  does not happen.

# Set Operations: Complement

## Definition

The **complement** of a set  $A \subseteq \Omega$ , denoted  $A^c$ , is the set of all elements in  $\Omega$  that are not in  $A$ .

When  $A$  is an event,  $A^c$  means that the event  $A$  does not happen.

Example:

$A = \{1, 3, 5\}$     “an odd roll”

# Set Operations: Complement

## Definition

The **complement** of a set  $A \subseteq \Omega$ , denoted  $A^c$ , is the set of all elements in  $\Omega$  that are not in  $A$ .

When  $A$  is an event,  $A^c$  means that the event  $A$  does not happen.

Example:

$A = \{1, 3, 5\}$     “an odd roll”

$A^c = \{2, 4, 6\}$     “an even roll”



# Set Operations: Difference

## Definition

The **difference** of a set  $A \subseteq \Omega$  and a set  $B \subseteq \Omega$ , denoted  $A - B$ , is the set of all elements in  $\Omega$  that are in  $A$  and are not in  $B$ .

Example:

$$A = \{3, 4, 5, 6\}$$

$$B = \{3, 5\}$$

$$A - B = \{4, 6\}$$

Note:  $A - B = A \cap B^c$

# DeMorgan's Law

Complement of union or intersection:

$$(A \cup B)^c = A^c \cap B^c$$

$$(A \cap B)^c = A^c \cup B^c$$

# DeMorgan's Law

Complement of union or intersection:

$$(A \cup B)^c = A^c \cap B^c$$

$$(A \cap B)^c = A^c \cup B^c$$

What is the English translation for both sides of the equations above?

# Exercises

Check whether the following statements are true or false.  
(Hint: you might use Venn diagrams.)

- ▶  $A - B \subseteq A$
- ▶  $(A - B)^c = A^c \cup B$
- ▶  $A \cup B \subseteq B$
- ▶  $(A \cup B) \cap C = (A \cap C) \cup (B \cap C)$

# Probability

## Definition

A **probability function** on a finite sample space  $\Omega$  assigns every event  $A \subseteq \Omega$  a number in  $[0, 1]$ , such that

1.  $P(\Omega) = 1$
2.  $P(A \cup B) = P(A) + P(B)$  when  $A \cap B = \emptyset$

$P(A)$  is the **probability** that event  $A$  occurs.

## Equally Likely Outcomes

The number of elements in a set  $A$  is denoted  $|A|$ .

# Equally Likely Outcomes

The number of elements in a set  $A$  is denoted  $|A|$ .

If  $\Omega$  has a finite number of elements, and each is equally likely, then the probability function is given by

$$P(A) = \frac{|A|}{|\Omega|}$$

# Equally Likely Outcomes

The number of elements in a set  $A$  is denoted  $|A|$ .

If  $\Omega$  has a finite number of elements, and each is equally likely, then the probability function is given by

$$P(A) = \frac{|A|}{|\Omega|}$$

Example: Rolling a 6-sided die



# Equally Likely Outcomes

The number of elements in a set  $A$  is denoted  $|A|$ .

If  $\Omega$  has a finite number of elements, and each is equally likely, then the probability function is given by

$$P(A) = \frac{|A|}{|\Omega|}$$

Example: Rolling a 6-sided die

▶  $P(\{1\}) = 1/6$

# Equally Likely Outcomes

The number of elements in a set  $A$  is denoted  $|A|$ .

If  $\Omega$  has a finite number of elements, and each is equally likely, then the probability function is given by

$$P(A) = \frac{|A|}{|\Omega|}$$

Example: Rolling a 6-sided die

- ▶  $P(\{1\}) = 1/6$
- ▶  $P(\{1, 2, 3\}) = 1/2$

# Repeated Experiments

If we do two runs of an experiment with sample space  $\Omega$ , then we get a new experiment with sample space

$$\Omega \times \Omega = \{(x, y) : x \in \Omega, y \in \Omega\}$$

# Repeated Experiments

If we do two runs of an experiment with sample space  $\Omega$ , then we get a new experiment with sample space

$$\Omega \times \Omega = \{(x, y) : x \in \Omega, y \in \Omega\}$$

The element  $(x, y) \in \Omega \times \Omega$  is called an **ordered pair**.

# Repeated Experiments

If we do two runs of an experiment with sample space  $\Omega$ , then we get a new experiment with sample space

$$\Omega \times \Omega = \{(x, y) : x \in \Omega, y \in \Omega\}$$

The element  $(x, y) \in \Omega \times \Omega$  is called an **ordered pair**.

Properties:

Order matters:  $(1, 2) \neq (2, 1)$

Repeats are possible:  $(1, 1) \in \mathbb{N} \times \mathbb{N}$

# More Repeats

Repeating an experiment  $n$  times gives the sample space

$$\begin{aligned}\Omega^n &= \Omega \times \cdots \times \Omega \quad (n \text{ times}) \\ &= \{(x_1, x_2, \dots, x_n) : x_i \in \Omega \text{ for all } i\}\end{aligned}$$

# More Repeats

Repeating an experiment  $n$  times gives the sample space

$$\begin{aligned}\Omega^n &= \Omega \times \cdots \times \Omega \quad (n \text{ times}) \\ &= \{(x_1, x_2, \dots, x_n) : x_i \in \Omega \text{ for all } i\}\end{aligned}$$

The element  $(x_1, x_2, \dots, x_n)$  is called an  **$n$ -tuple**.

# More Repeats

Repeating an experiment  $n$  times gives the sample space

$$\begin{aligned}\Omega^n &= \Omega \times \cdots \times \Omega \quad (n \text{ times}) \\ &= \{(x_1, x_2, \dots, x_n) : x_i \in \Omega \text{ for all } i\}\end{aligned}$$

The element  $(x_1, x_2, \dots, x_n)$  is called an  **$n$ -tuple**.

If  $|\Omega| = k$ , then  $|\Omega^n| = k^n$ .



# Probability Rules

# Probability Rules

Complement of an event  $A$ :

$$P(A^c) = 1 - P(A)$$

# Probability Rules

Complement of an event  $A$ :

$$P(A^c) = 1 - P(A)$$

Union of two overlapping events  $A \cap B \neq \emptyset$ :

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

# Exercise

You are picking a number out of a hat, which contains the numbers 1 through 100. What are the following events and their probabilities?

- ▶ The number has a single digit
- ▶ The number has two digits
- ▶ The number is a multiple of 4
- ▶ The number is not a multiple of 4
- ▶ The sum of the number's digits is 5

# Permutations

A **permutation** is an ordering of an  $n$ -tuple. For instance, the  $n$ -tuple  $(1, 2, 3)$  has the following permutations:

$(1, 2, 3), (1, 3, 2), (2, 1, 3)$   
 $(2, 3, 1), (3, 1, 2), (3, 2, 1)$

# Permutations

A **permutation** is an ordering of an  $n$ -tuple. For instance, the  $n$ -tuple  $(1, 2, 3)$  has the following permutations:

$$(1, 2, 3), (1, 3, 2), (2, 1, 3) \\ (2, 3, 1), (3, 1, 2), (3, 2, 1)$$

The number of unique orderings of an  $n$ -tuple is  **$n$  factorial**:

$$n! = n \times (n - 1) \times (n - 2) \times \cdots \times 2$$

# Permutations

A **permutation** is an ordering of an  $n$ -tuple. For instance, the  $n$ -tuple  $(1, 2, 3)$  has the following permutations:

$$(1, 2, 3), (1, 3, 2), (2, 1, 3) \\ (2, 3, 1), (3, 1, 2), (3, 2, 1)$$

The number of unique orderings of an  $n$ -tuple is  **$n$  factorial**:

$$n! = n \times (n - 1) \times (n - 2) \times \cdots \times 2$$

How many ways can you rearrange  $(1, 2, 3, 4)$ ?